

Struggling to reel in BV?

Release yourself from the discomfort of bacterial vaginosis (BV).

Solosec® (secnidazole) delivers a complete course of therapy in just **one oral dose.**¹

One packet. One dose. One time.

**ONE PACKET.
ONE DOSE.
ONE TIME.**



The entire contents of SOLOSEC packet should be sprinkled onto applesauce, yogurt, or pudding and consumed immediately (within max 30 minutes) **without chewing or crunching the granules.**¹

What is SOLOSEC?

SOLOSEC® (secnidazole) 2g oral granules is a prescription medicine used to treat bacterial vaginosis in adult women.

Select Important Safety Information

- You should not use SOLOSEC if you've had an allergic reaction to secnidazole, other ingredients of the formulation, or other nitroimidazole derivatives.

Please see complete Important Safety Information on the last page of this brochure.



Learn about bacterial vaginosis.

What is BV?

BV is a common vaginal infection. It occurs when there is an overgrowth of naturally-occurring bad bacteria that overwhelms your vagina's good bacteria.²

Women with BV don't always have symptoms, but you may notice common BV signs and symptoms like³:

- ▶ A thin white/gray vaginal discharge^{2,4}
- ▶ A noticeable fishy vaginal odor, especially after sex^{2,4}
- ▶ Itching in or around the vagina²

BV is the most common cause of vaginal symptoms, affecting

OVER 21 MILLION

A yellow silhouette map of the United States is positioned behind the text 'OVER 21 MILLION'.


women in the U.S. between the ages of 14 and 49 every year.^{2,4}

BV is not a yeast infection.

Some BV symptoms may seem similar to a yeast infection, but BV and yeast infections are **two different conditions**.^{2,5}

BV is bacterial, so it's treated by antibiotics which are **only** available by a prescription from a healthcare professional (HCP).³ Medications you can buy to treat yeast infections won't work for BV.⁵

Some factors can make you more likely to develop BV:





Talk to your doctor about BV.



If you think you have BV, speak to your HCP right away to get treated.³

You shouldn't wait for symptoms to go away or browse the web for home remedies. Self-treating with something you bought at the pharmacy isn't a good idea either, because over-the-counter medications aren't approved to treat BV.^{2,5}

Unless you know for sure what you have, there's no guarantee you can treat it properly.⁵



Untreated BV can negatively impact your feminine health.⁷

BV isn't "just a nuisance" or something to ignore. Leaving BV untreated can **increase your risk of:**

- ▶ Getting HIV, or transmitting HIV to your partner if you already have it⁷
- ▶ Getting sexually transmitted infections (STIs) like herpes, chlamydia, or gonorrhea⁷
- ▶ Developing pelvic inflammatory disease (PID), which can threaten your fertility⁷
- ▶ Experiencing preterm birth if you are pregnant⁷



BV often comes back.^{7,8}

Over 50% of women

will have BV again within a year of being successfully treated with antibiotics.⁸



If you're having **symptoms of BV again**, you should **speak to your HCP**.

What about antibiotic resistance?

Experts have studied whether BV comes back so often because of antibiotic resistance.⁹ There's evidence that some types of antibiotics can leave behind bacteria that are resistant to treatment.⁹

If you get diagnosed with BV again, your healthcare professional may select a different treatment for you than the one you were prescribed before.¹⁰



Learn about Solosec.

About Solosec® (secnidazole)


A full course of treatment for BV with **just one oral dose**¹



- ▶ Taken at any time of the day¹



- ▶ No intravaginal creams/gels or multiple days of pills¹¹⁻¹⁵



- ▶ No need for backup birth control if using a common birth control pill^{1*}

*Combination ethinyl estradiol and norethindrone



- ▶ No food or drink restrictions, including alcohol¹
– Laboratory studies show Solosec does not inhibit the enzyme that processes alcohol in your body¹

- ▶ Taken without regard to meals¹

Select Important Safety Information

- Take SOLOSEC exactly as your healthcare provider tells you to. Do not take SOLOSEC more often than it is prescribed.

Please see complete Important Safety Information on the last page of this brochure.



Treat yourself to a snack, and Solosec will begin working to treat your BV.¹

You've probably taken medications by mouth in the past, but they may have been pills, tablets, or capsules. You should know that Solosec is different.¹

ONE PACKET.
ONE DOSE.
ONE TIME.



The entire contents of SOLOSEC packet should be sprinkled onto applesauce, yogurt, or pudding and consumed immediately (within max 30 minutes) **without chewing or crunching the granules.**¹

Completing treatment with Solosec is as easy as 1,2,3:

1




Open the packet and **sprinkle** the entire contents of the packet onto applesauce, yogurt, or pudding¹

2



Stir and **mix**¹

3



Consume all of the mixture immediately, (within maximum 30 minutes). **DO NOT CHEW OR CRUNCH THE GRANULES**¹



Note: You can drink a glass of water after taking Solosec to help with swallowing, if needed. **Do not try to dissolve the medicine in water or any other liquid.**¹

BV**vibrant**

Get back to your life sooner.¹⁶

Here's what patients think.

Women responded to a recent survey about treatment with Solosec for BV*:

9 out of 10



- ▶ Were **satisfied, very satisfied, or extremely satisfied** with the result of their Solosec treatment^{17*}
- ▶ Would **recommend to a friend** with BV that they speak to their doctor about Solosec^{17*}

8 out of 10



- ▶ Found it **easy to take** their Solosec dose^{17*}
- ▶ **Would request Solosec again** if diagnosed with BV in the future^{17*}

* More than 4,000 Solosec kits were distributed to healthcare professionals, including a Solosec voucher, patient brochure, dosing card, and instruction card on how to participate in a brief voluntary survey. 1,888 Solosec vouchers were redeemed and 85 surveys were completed (N=85) in a nonscientific survey conducted by Lupin Pharmaceuticals. Numbers have been rounded.¹⁷



Learn what a real patient had to say about treatment with Solosec:



“ Solosec worked for me. My BV was taken care of with that one little packet. One packet, one time, and I was done with my treatment. ”

— *Lashera, Actual Solosec Patient**

*Individual results may vary

Solosec is generally well-tolerated¹

The most common side effects of Solosec include yeast infection, headache, nausea, altered taste, vomiting, diarrhea, abdominal pain, and vaginal itching.¹

Select Important Safety Information

- Before taking SOLOSEC, tell your healthcare provider about all of your medical conditions, including if you
 - Are pregnant or plan to become pregnant.
 - Are breastfeeding or plan to breastfeed. Breastfeeding is not recommended. You should not breastfeed for 96 hours (4 days) after taking SOLOSEC.

BV^{valued}

Save on Solosec.

There are multiple ways to access a co-pay card that may help you save money on your Solosec prescription*



If commercially insured, most eligible patients may pay **as little as \$25** for Solosec*:

- ▶ Click [here](#) to text a card to your mobile device, get a card emailed to you, or download a card onto your computer
- ▶ Alternatively, get a physical card from your HCP
- ▶ Using your mobile device, scan the QR code on this brochure below

* Per fill. Not valid for any person eligible for reimbursement of prescriptions, in whole or in part, by any state- or federally funded program, including Medicare, Medicaid, or Medicaid, or where prohibited, taxed or otherwise restricted by law. Cash-pay patients may pay as little as \$75. Eligibility requirements and terms and conditions apply. Click [here](#) for full details.



Select Important Safety Information

- You should not use SOLOSEC if you've had an allergic reaction to secnidazole, other ingredients of the formulation, or other nitroimidazole derivatives.

Please see complete Important Safety Information on the last page of this brochure.

Visit [solosec.com](https://www.solosec.com) for more information about BV and treatment with Solosec

Access Women's Health Resources

The more information you have, the better equipped you are to speak to your healthcare professional. Visit the following sites for information about women's health:

[American Sexual Health Association \(ASHA\)
ashasexualhealth.org](https://www.ashasexualhealth.org)

[Office on Women's Health \(OWH\)
womenshealth.gov](https://www.womenshealth.gov)

References: 1. SOLOSEC [prescribing information]. Baltimore, MD: Lupin Pharmaceuticals, Inc.; 2017. 2. Koumans EH, Sternberg M, Bruce C, et al. The prevalence of bacterial vaginosis in the United States, 2001-2004; associations with symptoms, sexual behaviors, and reproductive health. *Sex Transm Dis*. 2007;34(11):864-869. 3. Harvard Medical School. "Bacterial Vaginosis (Gardnerella Vaginitis)." *Harvard Health Publishing*. https://www.health.harvard.edu/a_to_z/bacterial-vaginosis-gardnerella-vaginitis-a-to-z. 4. Workowski KA, Bolan GA; CDC. Sexually Transmitted Diseases Treatment Guidelines, 2015 [published correction appears in *MMWR Recomm Rep*. 2015;64(33):924]. *MMWR Recomm Rep*. 2015;64(RR-03):1-137. 5. Orenstein BW. Types of vaginal infections. www.everydayhealth.com/news/vaginal-infection-types/. Accessed September 25, 2017. 6. Brookheart RT, Lewis WG, Peipert JF, Lewis AL, Allsworth JE. Association between obesity and bacterial vaginosis as assessed by Nugent Score. *Am J Obstet Gynecol*. 2019;220(5):476.e1-476.e11. 7. Lewis FM, Bernstein KT, Aral SO. Vaginal microbiome and its relationship to behavior, sexual health, and sexually transmitted diseases. *Obstet Gynecol*. 2017(4);129:643-654. doi:10.1097/AOG.0000000000001932. 8. Bradshaw CS, Morton AN, Hocking J, et al. High recurrence rates of bacterial vaginosis over the course of 12 months after oral metronidazole therapy and factors associated with recurrence. *J Infect Dis* 2006; 193:1478-89. 9. Bradshaw CS, Sobel JD. Current Treatment of Bacterial Vaginosis-Limitations and Need for Innovation. *J Infect Dis*. 2016;214 Suppl 1(Suppl 1):S14-S20. doi:10.1093/infdis/jiw159. 10. American College of Obstetricians and Gynecologists. ACOG Practice Bulletin: Clinical Management Guidelines for Obstetricians-Gynecologists, Number 215. *Obstet Gynecol* 2020;135(1):e1-e17. 11. Flagyl Extended Release [prescribing information]. Chicago, IL: G.D. Searle LLC; 2018. 12. Tinidazole Tablets [prescribing information]. Buffalo Grove, IL: G.D. PACK Pharmaceuticals, LLC; 2013. 13. Clindesse [prescribing information]. Allegan, MI: Perrigo Company, LLC; 2014. 14. Metrogel [prescribing information]. Sellersville, PA: Teva Pharmaceuticals USA; 2005. 15. Nuversa [prescribing information]. Madrid, Spain: Chemo Research SL; 2017. 16. Bilardi JE, Walker S, Temple-Smith M, et al. The burden of bacterial vaginosis: women's experience of the physical, emotional, sexual and social impact of living with recurrent bacterial vaginosis. *PLoS One*. 2013 Sep 11;8(9):e74378. doi:10.1371/journal.pone.0074378. 17. Data on file. Experience Program: Post Treatment Patient Survey data (May 2019-Sep 2019). Prepared September 2019. (LPSO-23506). Lupin Pharmaceuticals, Inc.



What is SOLOSEC?

SOLOSEC® (secnidazole) 2g oral granules is a prescription medicine used to treat bacterial vaginosis in adult women.

How should I take SOLOSEC?

SOLOSEC is a single-dose therapy for oral use. Use SOLOSEC by sprinkling an entire packet of SOLOSEC onto applesauce, yogurt, or pudding. The entire dose should be taken at once, and finished within 30 minutes. Avoid chewing or crunching the granules. SOLOSEC should not be taken by dissolving the granules in any liquid.

Important Safety Information

- You should not use SOLOSEC if you've had an allergic reaction to secnidazole, other ingredients of the formulation, or other nitroimidazole derivatives.
- Before taking SOLOSEC, tell your healthcare provider about all of your medical conditions, including if you
 - Are pregnant or plan to become pregnant.
 - Are breastfeeding or plan to breastfeed. Breastfeeding is not recommended. You should not breastfeed for 96 hours (4 days) after taking SOLOSEC.
- Vaginal yeast infections may occur with SOLOSEC and require an antifungal treatment.
- Take SOLOSEC exactly as your healthcare provider tells you to. Do not take SOLOSEC more often than it is prescribed.
- The most common side effects of SOLOSEC include yeast infection, headache, nausea, altered taste, vomiting, diarrhea, abdominal pain, and vaginal itching.

Call your doctor for medical advice on side effects. You are encouraged to report suspected side effects of prescription drugs to FDA at www.fda.gov/MedWatch also at 1-800-FDA-1088 or contact Lupin Pharmaceuticals, Inc. at 1-844-SOLOSEC (1-844-765-6732).

Please click [here](#) for full Prescribing Information, also available at solosec.com.